

Short-range correlations and entropy in ultracold atomic Fermi gases

Zhenhua Yu,^a Georg M. Bruun,^a and Gordon Baym^{a,b}

^a*The Niels Bohr International Academy, The Niels Bohr Institute,
Blegdamsvej 17, DK-2100 Copenhagen Ø, Denmark*

^b*Department of Physics, University of Illinois, 1110 W. Green Street, Urbana, IL 61801*

(Dated: May 12, 2009)

We relate short-range correlations in ultracold atomic Fermi gases to the entropy of the system over the entire temperature, T , vs. coupling strength, $-1/k_F a$, plane. In the low temperature limit the entropy is dominated by phonon excitations and the correlations increase as T^4 . In the BEC limit, we calculate a boson model within the Bogoliubov approximation to show explicitly how phonons enhance the fermion correlations. In the high temperature limit, we show from the virial expansion that the correlations decrease as $1/T$. The correlations therefore reach a maximum at a finite temperature. We infer the general structure of the isentropes of the Fermi gas in the $T, -1/k_F a$ plane, and the temperature dependence of the correlations in the unitary, BEC, and BCS limits. Our results compare well with measurements of the correlations via photoassociation experiments at higher temperatures.

PACS numbers:

I. INTRODUCTION

The thermodynamics of an ultracold atomic gas interacting via short range interactions is encoded in the short range two-body correlations of the particles. Such connections between short range correlations and physical properties of systems with short range interactions have been long recognized, e.g., Ref. [1], and have been employed recently to study the rf spectra of paired Fermi gases [2, 3, 4]. Tan has derived a number of remarkable relations between the properties of ultracold atomic gases, e.g., the ground state energy and their short range correlations [5] (see also [6]). These links open new perspectives to examine the many particle physics of systems with short range interactions.

By using Feshbach resonances to tune the scattering length a , one can study atomic Fermi gases through the crossover from a weakly coupled BCS superfluid to a weakly coupled Bose-Einstein condensate (BEC) of molecules. In the strongly correlated unitarity regime, $|a| \rightarrow \infty$, the many-body correlations are highly non-trivial and one has to resort to Monte-Carlo calculations to obtain controlled results [7, 8]. At non-zero temperatures T , information on the correlations is even more limited. We study here how knowledge of the free energy and entropy can illuminate the nature of the finite temperature short range two particle correlations. The two-body correlation function in a two component Fermi gas has the general structure

$$\langle \psi_1^\dagger(\mathbf{r}') \psi_2^\dagger(0) \psi_2(0) \psi_1(\mathbf{r}) \rangle = \sum_i \gamma_i \phi_i(\mathbf{r}) \phi_i^*(\mathbf{r}'), \quad (1)$$

as one sees by regarding the correlation function as a Hermitean operator in $\mathbf{r}, \mathbf{r}'$. For $r \ll d$, the interparticle spacing, the functions ϕ_i , essentially s-wave Jastrow factors in the many-body wavefunction at short interparticle distance, are determined by two-body physics, and outside the range of the potential, r_0 , at temper-

atures of interest, are $\sim \sin(kr + \delta)/r = \sin \delta \chi(r)/r$, where $\chi(r) = 1 - r/a$, δ is the s-wave phase shift, and $\cot \delta = -1/ka$, where a is the s-wave scattering length. Thus

$$\langle \psi_1^\dagger(\mathbf{r}) \psi_2^\dagger(0) \psi_2(0) \psi_1(\mathbf{r}) \rangle = C \left(\frac{\chi(r)}{r} \right)^2 = C \left(\frac{1}{r} - \frac{1}{a} \right)^2. \quad (2)$$

The correlation (or contact) strength C , which is determined by many-body physics, can be measured directly in photoassociation experiments [9, 10].

Here we examine the temperature dependence of the correlations by calculating $C(T)$ from the finite temperature thermodynamics. Quite generally, the short range correlations are related to the free energy density $f = \text{Tr} e^{-H/T} / \Omega$ (where Ω is the volume of the system) by [4]:

$$C = -\frac{m}{4\pi} \frac{\partial f}{\partial a^{-1}}. \quad (3)$$

Differentiating with respect to the temperature, T , we find

$$\frac{\partial C}{\partial T} = -\frac{m}{4\pi} \frac{\partial}{\partial T} \frac{\partial f}{\partial a^{-1}} = \frac{m}{4\pi} \frac{\partial s}{\partial a^{-1}}, \quad (4)$$

thus linking the variation of C with temperature to the coupling constant dependence of the entropy density s . The connection is illustrated below in Fig. 1 which shows the temperature dependence of C , and the related Fig. 2, which shows the isentropes in the $T, -1/k_F a$ plane.

We focus on the unitarity, BEC, and BCS limits and provide controlled results for C for both high and low T . Using the virial expansion, we show that the correlations decrease with $C \propto n^2/T$ in the high temperature limit, where n is the density. From the relation (4) between the T dependence of C and the entropy density, we show that $C(T)$ increases with temperature as T^4 for low T , owing

to the phonon contribution to the entropy. In the BEC limit, we assume a bosonic model to relate the bosonic and fermionic correlations; a calculation within the Bogoliubov approximation illustrates how phonons enhance the fermion correlations at low temperature. Combining the behavior of C in the high and low T limits, we infer that C reaches a maximum value at a temperature $T_{\max} > 0$. This intriguing behavior of $C(T)$ is detectable in photoassociation experiments near unitarity.

II. CORRELATION FUNCTION AND FREE ENERGY

We consider a homogenous two-component Fermi gas described by the Hamiltonian ($\hbar = 1$ throughout)

$$H = \int d^3r \sum_{i=1,2} \frac{1}{2m} \nabla \psi_i^\dagger(\mathbf{r}) \cdot \nabla \psi_i(\mathbf{r}) + \int d^3r \int d^3r' V(|\mathbf{r} - \mathbf{r}'|) \psi_1^\dagger(\mathbf{r}) \psi_2^\dagger(\mathbf{r}') \psi_2(\mathbf{r}') \psi_1(\mathbf{r}), \quad (5)$$

where the ψ_i are the field operators for fermions in internal state $|i\rangle$. We take the range r_0 of the interaction $V(r)$ to be short compared to the interparticle spacing d .

We first present an elementary derivation of the relation (3) between the two-body correlation function and the free energy density, using the device of scaling the potential $V(r)$ by a factor λ in (5); $V(r) \rightarrow \lambda V(r)$, letting $\lambda \rightarrow 1$ in the end. The correlation function for given λ has the structure of (2) with χ , C and a functions of λ . For $r \ll d$, the function χ_λ satisfies the low energy limit of the two-body Schrödinger equation [4, 11]

$$\left(-\frac{1}{m} \frac{d^2}{dr^2} + \lambda V(r) \right) \chi_\lambda(r) = 0. \quad (6)$$

At short distances, with $r > r_0$, $\chi_\lambda(r) = 1 - r/a_\lambda$; for convenience, we choose the normalization constant to be unity. Differentiating (6) with respect to λ , multiplying by χ_λ , integrating with respect to r from 0 to r_c , and eliminating the term $\int dr \chi_\lambda V \partial \chi_\lambda / \partial \lambda$ using (6), we find

$$\int_0^\infty dr V(r) \chi_\lambda^2(r) \frac{\partial \lambda}{\partial a_\lambda^{-1}} = -\frac{1}{m}, \quad (7)$$

where we use vanishing of the s-wave function χ_λ at the origin, and extend the upper bound of the integral from r_c to infinity since $V(r)$ is nonzero only for $r < r_0$. On the other hand, from the Feynman-Hellmann theorem, the variation of the free energy density with coupling

strength is

$$\begin{aligned} \frac{\partial f}{\partial a_\lambda^{-1}} &= \frac{\partial f}{\partial \lambda} \frac{\partial \lambda}{\partial a_\lambda^{-1}} = \frac{1}{\Omega} \left\langle \frac{\partial H}{\partial \lambda} \right\rangle \frac{\partial \lambda}{\partial a_\lambda^{-1}} \\ &= \int d^3r V(r) \langle \psi_1^\dagger(\mathbf{r}) \psi_2^\dagger(0) \psi_2(0) \psi_1(\mathbf{r}) \rangle_\lambda \frac{\partial \lambda}{\partial a_\lambda^{-1}} \\ &= 4\pi C_\lambda \int_0^\infty dr V(r) \chi_\lambda^2(r) \frac{\partial \lambda}{\partial a_\lambda^{-1}}. \end{aligned} \quad (8)$$

Combining (7) and (8), and setting $\lambda = 1$, we arrive at the desired relation (3).

We review the application of (3) to a homogeneous equally populated two-component Fermi gas of volume Ω , at $T = 0$ [10]. The ground state energy per particle $E/2N$ can be written as $(3/5)(1 + \beta)E_F$, where β is a function of the dimensionless quantity $\xi = -1/k_F a$. The Fermi momentum k_F is defined in terms of the single component density $n = N/\Omega$ by $n = k_F^3/6\pi^2$, and the Fermi energy by $E_F = k_F^2/2m$. Equation (3) implies that $C(T = 0) = (k_F^4/40\pi^3) \partial \beta / \partial \xi$. The function β has been calculated by quantum Monte Carlo simulations near unitarity ($1/a \rightarrow 0$); at unitarity $\partial \beta / \partial \xi \approx 0.9$ [7], a result which agrees well with the value $C = 2.7(k_F^4/36\pi^4)$ extracted directly from the correlation function calculated by quantum Monte Carlo methods [8]. In the BCS limit, $-1/k_F a \gg 1$, a perturbative expansion in $k_F a$ gives [12, 13, 14]

$$\frac{E}{2N} = \frac{3}{5} E_F \left[1 + \frac{10}{9\pi} k_F a + \dots \right]. \quad (9)$$

Corrections to E due to pairing are exponentially small in this limit. Keeping the leading term yields $C = a^2 n^2$. In the BEC limit, $1/k_F a \gg 1$, fermions form molecules and [12, 13]

$$\frac{E}{2N} = \frac{E_b}{2} + \frac{E_F k_F a_m}{6\pi} \left(1 + \frac{128}{15\sqrt{6}\pi^3} (k_F a_m)^{3/2} + \dots \right), \quad (10)$$

with $E_b = -1/ma^2$ the molecular binding energy, and a_m the scattering length between molecules. The second term is the Hartree-Fock mean field energy and the third the nonperturbative Lee-Yang correction [13]. Few body calculations give $a_m = 0.6a$ [15]. In the BEC limit, $C = n/2\pi a$ to leading order. The divergent behavior, as $a \rightarrow 0$, arises from the normalization of the molecular wave function, $\chi/r \sim e^{-r/a}/ra^{1/2}$.

III. LOW TEMPERATURE

We now consider the temperature dependence of C for T well below the superfluid transition temperature T_c , in which regime phonon excitations dominate. The contribution of the phonons to the entropy density is

$$s_{\text{phonon}} = \frac{2\pi^2}{45} \left(\frac{T}{c_s} \right)^3, \quad (11)$$

where the zero temperature sound velocity c_s is given by $mc_s^2 = n\partial\mu/\partial n$, with μ is the chemical potential. The change of C to leading order for $T \ll T_F = E_F$ ($k_B = 1$) is thus

$$\delta C = C(T) - C(0) = -\frac{\pi m}{120} \left(\frac{T}{c_s}\right)^4 \frac{\partial c_s}{\partial a^{-1}}. \quad (12)$$

The zero temperature sound velocity appears to increase monotonically from the BEC side to the BCS side [7, 16, 17], so that the short range pair correlations as parametrized via C increase from zero temperature as T^4 . At $T = 0$, $\mu = E_F(1 + \beta - \beta'\xi/5)$, so that

$$\left(\frac{c_s}{v_F/\sqrt{3}}\right)^2 = 1 + \beta - \frac{3}{5}\beta'\xi + \frac{1}{10}\beta''\xi^2, \quad (13)$$

with $\beta' = \partial\beta/\partial\xi$, etc. Recent measurements of c_s are in good agreement with (13) combined with Monte Carlo results for β around unitarity [21].

Equation (9) implies that in the BCS limit, $c_s^2 = (v_F^2/3)(1 + 2k_F a/\pi)$, and thus at low T ,

$$C(T) = n^2 a^2 \left[1 + \frac{9\sqrt{3}\pi^4}{160} \left(\frac{T}{T_F}\right)^4 \right]. \quad (14)$$

For $T_c < T \ll T_F$, the weakly attractive Fermi gas can be described by Landau Fermi liquid theory. The entropy density is given by [25] $s_f = \frac{1}{3}m^*k_F T$ with $m^* = m[1 + 8(7\log 2 - 1)(k_F a)^2/15\pi^2]$ the effective mass of the quasiparticles [26]. From (4), we then have the temperature dependence

$$\delta C(T)/n^2 a^2 = -\frac{\pi}{5}(7\log 2 - 1)k_F a \left(\frac{T}{T_F}\right)^2 > 0. \quad (15)$$

In the BEC limit, the Bogoliubov sound speed is $c_s = \sqrt{ng_m/M}$ with $g_m = 4\pi a_m/M$ and $M = 2m$ [22]. Thus

$$C(T) = \frac{n}{2\pi a} \left[1 + \left(\frac{\partial a_m}{\partial a}\right) \frac{(k_F a)^3}{12\pi} \left(1 + \frac{64}{3\sqrt{6}\pi^3} (k_F a_m)^{3/2} + \frac{(486\pi^{13})^{1/2}}{40(k_F a)^{5/2}} \left(\frac{T}{T_F}\right)^4 \right) \right]. \quad (16)$$

At unitarity,

$$C(T) = \frac{k_F^4 \beta'}{40\pi^3} \left[1 + \frac{\sqrt{3}\pi^4}{80(1 + \beta)^{5/2}} \left(\frac{T}{T_F}\right)^4 \right]. \quad (17)$$

IV. CORRELATIONS IN THE BEC LIMIT

The universal behavior $\delta C \sim T^4$ as $T \rightarrow 0$ shows that thermally excited phonons enhance the fermion pair correlations. In the BEC limit, the enhancement of the correlations can be understood directly in terms of a gas of molecules interacting with a short range potential with a

scattering length $a_m > 0$. The boson correlation function is similar in structure to that for fermions:

$$\lim_{r \rightarrow 0} \langle \phi^\dagger(\mathbf{r}) \phi^\dagger(0) \phi(0) \phi(\mathbf{r}) \rangle = C_m \left(\frac{1}{r} - \frac{1}{a_m} \right)^2, \quad (18)$$

where the $\phi(\mathbf{r})$ are the bosonic field operators. As for fermions,

$$C_m = -\frac{M}{2\pi} \frac{\partial f_m}{\partial a_m^{-1}}, \quad (19)$$

with f_m the free energy density for the bosonic molecules. The factor of 2 difference from (3) is due to the factor 1/2 in the interaction energy for identical bosons written in terms of the ϕ .

Since the low temperature physics described in terms of the fundamental fermions or bosonic molecules must be the same, the leading temperature variation $\delta f = f(T) - f(0)$ is equal to $\delta f_m = f_m(T) - f_m(0)$. Therefore

$$\begin{aligned} \delta C &= -\frac{m}{4\pi} \frac{\partial(\delta f)}{\partial a^{-1}} = -\frac{m}{4\pi} \frac{\partial(\delta f_m)}{\partial a_m^{-1}} \frac{\partial a_m^{-1}}{\partial a^{-1}} \\ &= \frac{m}{2M} \frac{\partial a_m^{-1}}{\partial a^{-1}} \delta C_m, \end{aligned} \quad (20)$$

with $\delta C_m = C_m(T) - C_m(0)$. We calculate the boson correlation function within the Bogoliubov approximation using a pseudopotential $V_m(r) = g_m \delta(\mathbf{r})$. We divide the field operator $\phi(\mathbf{r}) = \phi_0 + \delta\phi(\mathbf{r})$ into a condensate part ϕ_0 and a fluctuation $\delta\phi$, assumed to be small. To second order in the fluctuations, at small r ,

$$\begin{aligned} \langle \phi^\dagger(\mathbf{r}) \phi^\dagger(0) \phi(0) \phi(\mathbf{r}) \rangle \\ = n_0^2 + 2n_0[2\langle \delta\phi^\dagger(0) \delta\phi(0) \rangle + \langle \delta\phi^\dagger(\mathbf{r}) \delta\phi^\dagger(0) \rangle], \end{aligned} \quad (21)$$

where

$$\langle \delta\phi(0)^\dagger \delta\phi(0) \rangle = \int \frac{d^3 k}{(2\pi)^3} [v_k^2 + (u_k^2 + v_k^2) \langle \alpha_k^\dagger \alpha_k \rangle], \quad (22)$$

$$\begin{aligned} \lim_{r \rightarrow 0} \langle \delta\phi^\dagger(\mathbf{r}) \delta\phi^\dagger(0) \rangle &= -\int \frac{d^3 k}{(2\pi)^3} u_k v_k [e^{i\mathbf{k} \cdot \mathbf{r}} + 2\langle \alpha_k^\dagger \alpha_k \rangle] \\ &= -\frac{n_0 a_m}{r} - \int \frac{d^3 k}{(2\pi)^3} \left(u_k v_k - \frac{M g_m n_0}{k^2} + 2u_k v_k \langle \alpha_k^\dagger \alpha_k \rangle \right); \end{aligned} \quad (23)$$

here $v_k^2 = (\xi_k/E_k - 1)/2$, $u_k^2 = (\xi_k/E_k + 1)/2$, $\xi_k = k^2/2M + g_m n_0$, and the α_k annihilate phonons with energy dispersion $E_k = \sqrt{\xi_k^2 - (g_m n_0)^2}$. The $1/r$ divergence in $\langle \delta\phi^\dagger(\mathbf{r}) \delta\phi^\dagger(0) \rangle$ corresponds to the cross term in (18). The $\sim 1/r^2$ in (18) appears from the higher order term $\lim_{r \rightarrow 0} \langle \delta\phi^\dagger(\mathbf{r}) \delta\phi^\dagger(0) \rangle \langle \delta\phi(\mathbf{r}) \delta\phi(0) \rangle = \lim_{r \rightarrow 0} |\langle \delta\phi(\mathbf{r}) \delta\phi(0) \rangle|^2 + \dots$, not included in (21).

The lowest order Bogoliubov approximation produces the correlation function (18) to leading order in $(na_m^3)^{1/2}$. In (21), the r independent term is reliable to order $(na_m^3)^{1/2}$, the term proportional to $1/n^{1/3}r$ to order

$(na_m^3)^{1/3}$, and the term proportional to $1/(n^{1/3}r)^2$ to order $(na_m^3)^{2/3}$. To derive the full $C_m(1 - a_m/r)^2$ structure to higher order in $(na_m^3)^{1/2}$ requires calculating terms beyond the simple ones in (21), a task we defer. To extract C_m from the r independent term in (21) we write $n_0 = n - \langle \delta\phi(0)^\dagger \delta\phi(0) \rangle$. At $T = 0$, $C_m/(na_m)^2 = 1 + (64/3)(na_m^3/\pi)^{1/2}$, which agrees with the calculation of C_m from (19) using the ground state energy of weakly interacting bosons [cf. (10)]. The thermally induced change δC_m is

$$\begin{aligned} \frac{\delta C_m}{2na_m^2} &= \int \frac{d^3k}{(2\pi)^3} (u_k^2 + v_k^2 - 2u_k v_k) \langle \alpha_k^\dagger \alpha_k \rangle \\ &= \int \frac{d^3k}{(2\pi)^3} \frac{k^2}{2E_k} \frac{1}{e^{E_k/T} - 1} = \frac{\pi^2 T^4}{60 c_s^5}, \end{aligned} \quad (24)$$

where we consider only the phonon contribution ($E_k = c_s k$). Substituting (24) into (20), we obtain the same result as in (16).

The T^4 dependence can be understood in terms of the effect of thermal sound waves on the eigenstates, $\phi_i(r)$, of the two particle density matrix (1). At distances beyond the interparticle spacing, Eq. (6) for the ϕ (with $\lambda = 1$) contains potential terms from the mean field $g_m n$ and its fluctuations. At fixed normalization, the latter change the magnitude of $\phi_i(r)$ at short distances by terms $\sim \langle (\delta n)^2 \rangle \sim T^4$, which translates into the T^4 dependence of C at low temperature.

V. HIGH TEMPERATURE

In the high temperature limit, $T \gg T_F$, where the fugacity $z = e^{\mu/T}$ is small, the free energy can be calculated via a virial expansion. To second order in z , the partition function for a two-component Fermi gas is given by [23, 24]

$$\log(Z/Z_0) = \frac{2^{3/2}\Omega}{\lambda^3} z^2 b_2, \quad (25)$$

where

$$b_2 = \sum_i e^{E_b^i/T} + \int_0^\infty \frac{dk}{\pi} \frac{d\delta(k)}{dk} e^{-k^2/mT} \quad (26)$$

is the second virial coefficient. Here the E_b^i are the bound energies of the two-body attractive interaction $V(r)$, $\delta(k)$ is the s-wave scattering phase shift, and $\lambda = (2\pi/mT)^{1/2}$ is the thermal wavelength. In the BEC regime, we consider only the relevant bound state with $E_b = -1/ma^2$. In the regime $T \ll 1/mr_0^2$, the values of k entering (26) are $\ll 1/r_0$, so that $\cot \delta(k) = -1/ka$; hence $d\delta(k)/dk = -a/(ka^2 + 1)$, and

$$b_2 = \left(\frac{1}{2} + \frac{1}{\sqrt{\pi}} \int_0^{\lambda/\sqrt{2\pi}a} e^{-t^2} dt \right) e^{\lambda^2/2\pi a^2}, \quad (27)$$

a dimensionless function of λ/a . Thus

$$C = \frac{mkT}{4\pi\Omega} \frac{\partial \log Z}{\partial a^{-1}} = \sqrt{2} n^2 \lambda^2 \frac{\partial b_2}{\partial(\lambda/a)}, \quad (28)$$

where we replace z by the number density $n = (T/\Omega)(\partial \log Z/\partial \mu) = (z/\lambda^3)(1 + 2^{3/2} z b_2)$.

For $T \gg T_a \equiv 1/ma^2$ ($a \gg \lambda$), the system is a weakly interacting Fermi gas for all coupling strengths. From (27), $\partial b_2/\partial(\lambda/a) = 1/\sqrt{2}\pi$, and thus

$$C(T) = \frac{n^2 \lambda^2}{\pi} = \frac{2n^2}{mT}. \quad (29)$$

Equation (29) combined with (12) leads to the important result that the correlation C has a maximum at a *nonzero* temperature T_{max} .

In the BCS limit, for $T_F \ll T \ll T_a$, $b_2 = -a/\sqrt{2}\lambda$ and $C(T) = n^2 a^2$; this is simply the Hartree term in the low T result (14) persisting to higher temperatures. From (14), (15) and (29), we expect that $T_{max} \sim T_F$ and that the relative size of the maximum, $[C(T_{max}) - C(0)]/C(0) \sim |k_F a|$, is small.

In the BEC limit, when the temperature is lowered to $T \sim T_a$ ($\lambda \sim a$), the bound state starts to dominate in (27) and $b_2 \sim \exp(\lambda^2/2\pi a^2)$. The correlation C increases from $\sim n^2 \lambda^2$ to $\sim n^2 \lambda^3 \exp(\lambda^2/2\pi a^2)/a$. The virial expansion for the Fermi gas breaks down when $b_2 z \sim 1$, which happens at $T \sim T_a/\log(1/k_F a)$. At this point the number of the bosonic molecules becomes significant. Within the regime $T_F \ll T \ll T_a/\log(1/k_F a)$, a virial expansion can be carried out for the bosonic molecules. The calculation proceeds as for the fermions except that there is no bound state for the molecules and one has to add the binding energy $-n/ma^2$ to the free energy density, yielding

$$C(T) = \frac{n}{2\pi a} \left[1 + \frac{(k_F a)^3}{6\pi} \left(\frac{\partial a_m}{\partial a} \right) \right]. \quad (30)$$

The second term in (30) comes from the mean field energy for a normal weakly interacting Bose gas, which is twice that of a condensed Bose gas [cf. (16)]. From (16), (29) and (30), we expect that C reaches a maximum at $T_{max} \sim T_c \sim T_F$, and that $[C(T_{max}) - C(0)]/C(0) \ll 1$.

In the unitarity limit, (17) and (29) indicate that C has a maximum at $T_{max} \sim T_F$ with $[C(T_{max}) - C(0)]/C(0) \sim 1$. The pronounced maximum in the correlations in the strong coupling limit indicates the importance of phonons there.

Figure 1 sketches $C(T)$ in the BCS, unitarity, and BEC regimes. The solid curves are interpolations between the low T results (14)-(17) and the virial expansion results (28)-(30) for $T \gg T_F$. Note that C is continuous at T_c since the superfluid transition is second order.

The link between the entropy and the short range correlations through (4) is illustrated in Fig. (2), where we sketch the isentropes in the BEC-BCS crossover regime, as well as the superfluid transition temperature




FIG. 1: (color online) The correlation strength $C(T)$ in the BCS (a), unitary (b), and BEC (c) limits. The dashed lines at high T are the virial results, and the dashed line at low T in (b) is from (17). The maximum in $C(T)$ for $T = T_{\max}$ in the BCS and BEC limits is too small to be visible here.

T_c [18, 19, 20]. It follows from (4) and the virial expansion result (29) that the isentropes has a negative slope in the $T, -1/k_F a$ plane for high T . Likewise, the increase in $C(T)$ given by (14)–(17) means that the isentropes has a positive slope for low T . The correlation strength $C(T)$ has an extremum where the slope of the isentropes as a function of $-1/k_F a$ vanishes. Additional information is obtained from a recent experiment, where a Fermi gas was prepared in the BCS limit with initial energy E_i and entropy S_i and then adiabatically tuned to the unitarity point where the final energy E_f was measured [21]. The temperature was deduced from $T = \partial E / \partial S$. Within the experimental regime the final temperatures T_f at unitarity were generally higher than the initial temperatures T_i in the BCS limit [21]. However, for the isentrope ending at $T_f \approx 0.2T_F$, $T_i \approx T_f$, which indicates that this isentrope (as well as neighboring ones) first bends upwards and then downwards from the unitarity region to the BCS limit. Well below the BCS transition, where phonons dominate the entropy, the isentropes have positive slope, not visible on the scale of Fig. (2).

VI. PHOTOASSOCIATION EXPERIMENT

Photoassociation experiments provide a direct measure of the correlation strength [9, 10]. In a recent experiment, ^6Li atoms were trapped in the lowest two hyperfine states $|1\rangle$ and $|2\rangle$. Then the bare closed channel spin-singlet molecular state $|\psi_{v=38}\rangle$ associated with the 834G Feshbach resonance in the 1-2 channel was excited by a laser field $\mathbf{E}_L$ to a spin-singlet molecular state $|\psi_{v'=68}\rangle$, with linewidth $\gamma = (2\pi)11.7$ MHz. The excited molecules were lost from the trap and the remaining atoms counted for




FIG. 2: (color online) Isentropes of a balanced two-component Fermi gas, from which one can infer the temperature dependence of $C(T)$.

various durations of the laser pulse [9]. By Fermi's golden rule, the local loss rate of the number of atoms, $2n$, in the trap is [10]

$$\Gamma = -2 \frac{dn(t)}{dt} = 2 \frac{\Omega_R^2}{\gamma} n_b, \quad (31)$$

where $\Omega_R = \langle \psi_{v'=68} | \mathbf{E}_L \cdot \mathbf{d} | \psi_{v=38} \rangle$ is the Rabi frequency, $\mathbf{d}$ is the atomic dipole operator, and n_b the local density of molecules in the closed channel. The density of closed channel molecules n_b is related to the correlation function between the states $|1\rangle$ and $|2\rangle$ in the open channel by [10]

$$n_b = \frac{4\pi a_{bg} C}{m\mu_b \Delta B} \left(\frac{1}{a_{bg}} - \frac{1}{a} \right)^2, \quad (32)$$

where a_{bg} is the background scattering length, the molecular magnetic moment μ_b is twice the Bohr magneton μ_B , and ΔB is the width of the Feshbach resonance.

In a homogeneous gas in the high temperature limit, we find from (29), (31), and (32) that $\Gamma \sim n^2/T$. In a trap, the local density approximation gives $n(\mathbf{r}) = N(\bar{\omega}/\lambda T)^3 \exp[-\beta V(\mathbf{r})]$ with $V(\mathbf{r}) = (m/2)(\omega_x^2 x^2 + \omega_y^2 y^2 + \omega_z^2 z^2)/2$ and $\bar{\omega} = (\omega_x \omega_y \omega_z)^{1/3}$. The average of C over the trap, and thus the average rate $\bar{\Gamma}$, is $\propto \bar{\omega}^3 N^2 / T^{5/2}$, and Eq. (31) has the solution

$$\frac{N(t)}{N(0)} = \left(1 + \frac{\Omega_R^2 N_b t}{\gamma N} \right)^{-1}. \quad (33)$$

Analogous expressions for the $T = 0$ trap-averaged loss rate were given in [10].

In Fig. 3, we compare the high T result (33) with the experimental data obtained on the BCS side of the resonance [9]. Initially the temperature is below T_f , where




FIG. 3: (color online) Loss of ${}^6\text{Li}$ atoms in the trap through photoassociation vs. the probe duration at $B = 865$ G. The solid curve is the numerical evaluation of (33). The circles are the $T = 0.75T_F$ experimental data from Ref. [9].

(31) underestimates the loss; however, owing to depletion the Fermi temperature falls below T , and (33) provides reasonable agreement with the data – with no fitting parameters.

Acknowledgements

This work grew out of the Niels Bohr International Academy Summer Institute on Cold Atoms and Quark-Gluon Plasmas. We are particularly indebted to Randy Hulet for discussions there of the Rice photoassociation experiments and providing us with the experimental data. This research was supported in part by NSF Grant PHY07-01611.

-
- [1] M. Fisher, Phil. Mag. **7** 1731 (1962); M. Fisher and J. Langer, Phys. Rev. Lett. **20** 655 (1968).
 - [2] G. Baym, C. Pethick, Z. Yu and M. Zwierlein, Phys. Rev. Lett. **99**, 190407 (2007).
 - [3] M. Punk and W. Zwerger, Phys. Rev. Lett. **99**, 170404 (2007).
 - [4] S. Zhang and A.J. Leggett, Phys. Rev. A **79**, 023601 (2009).
 - [5] S. Tan, Ann. of Phys. **323** 2952 (2008); *ibid* **323** 2971 (2008).
 - [6] E. Braaten and L. Platter, Phys. Rev. Lett. **100** 205301 (2008).
 - [7] G.E. Astrakharchik, J. Boronat, J. Casulleras, and S. Giorgini, Phys. Rev. Lett. **93**, 200404 (2004).
 - [8] C. Lobo, I. Carusotto, S. Giorgini, A. Recati, and S. Stringari, Phys. Rev. Lett. **97** 100405 (2006).
 - [9] G. B. Partridge, K. E. Strecker, R. I. Kamar, M. W. Jack, and R. G. Hulet, Phys. Rev. Lett. **95**, 020404 (2005).
 - [10] F. Werner, L. Tarruell, and Y. Castin, Euro. Phys. J. B (online, 2009), DOI: 10.1140/epjb/e2009-00040-8; arXiv:0807.0078.
 - [11] A.J. Leggett, *Quantum liquids: Bose condensation and Cooper pairing in condensed-matter systems*, (Oxford Univ. Press, Oxford, 2006).
 - [12] K. Huang and C.N. Yang, Phys. Rev. **105**, 767 (1957).
 - [13] T.D. Lee and C.N. Yang, Phys. Rev. **105**, 1119 (1957).
 - [14] V. M. Galitskii, JETP, **34**:151 (1958), 1011 [Engl. transl. Soviet Phys. JETP, **7**:104 (1958), 698].
 - [15] D.S. Petrov, C. Salomon, and G.V. Shlyapnikov, Phys. Rev. Lett. **93**, 090404 (2004); I.V. Brodsky, M.Y. Kagan, A.V. Klapptsov, R. Combescot, X. Leyronas, Phys. Rev. A **73**, 032724 (2006).
 - [16] J. Joseph, B. Clancy, L. Luo, J. Kinast, A. Turlapov, and J.E. Thomas, Phys. Rev. Lett. **98**, 170401 (2007).
 - [17] R. Combescot, M. Y. Kagan, and S. Stringari, Phys. Rev. A **74**, 042717 (2006).
 - [18] G. Baym, J.-P. Blaizot, M. Holzmann, F. Laloë, and D. Vautherin, Phys. Rev. Lett. **83**, 1703 (1999).
 - [19] E. Burovski, E. Kozik, N. Prokofev, B. Svistunov, and M. Troyer, Phys. Rev. Lett. **101**, 090402 (2008).
 - [20] L.P. Gor'kov and T.K. Melik-Barkhudarov, Sov. Phys. JETP **13**, 1081 (1961).
 - [21] L. Luo and J.E. Thomas, J. Low Temp. Phys. **154**, 1 (2009).
 - [22] C. J. Pethick and H. Smith, *Bose-Einstein condensation in dilute gases*, 2nd ed. (Cambridge Univ. Press, Cambridge, 2008).
 - [23] L.D. Landau and E.M. Lifshitz, *Statistical physics I* (Pergamon Press, Oxford, 1980).
 - [24] T-L Ho and E.J. Mueller, Phys. Rev. Lett. **92**, 160404 (2004).
 - [25] G. Baym and C.J. Pethick, *Landau Fermi-liquid theory* (John Wiley & Sons, Inc, New York, 1991).
 - [26] E.M. Lifshitz and L.P. Pitaevskii, *Statistical physics II* (Pergamon Press, Oxford, 1980).